



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ON A NEW FORM OF *OCEANODROMA* INHABITING
SAN BENITO ISLAND, OFF THE COAST OF
LOWER CALIFORNIA.

BY HANS GRAF VON BERLEPSCH.

***Oceanodroma monorhis chapmani* Berl.**

O. O. monorhis dictæ valde affinis sed alis multo, cauda tarsisque paulo brevioribus, rostro longiore, neonon corpore supra nigrescentiore, minus griseo lavato, distinguenda.

Al. 149-147 caud. 77-76 culm. $28\frac{3}{4}$ - $28\frac{1}{4}$ tars. $22\frac{1}{2}$ -22 mm.

Habitat in insula San Benito dicta, maris pacifici Californiæ inferiori proxima.

Typus in Mus. H. v. B. (♂ San Benito Isld.).

In the year 1899 I received through Mr. H. W. Marsden of Clifton, Bristol, a number of rare North American birds, which he had to sell in commission for Mr. C. K. Worthen of Warsaw, Ill., whose firm name is to be found printed on the labels. From this consignment I picked out several specimens of Petrels belonging to the genus *Oceanodroma*, collected in the San Benito and Guadalupe Islands off the coast of Lower California. They bear labels in a handwriting I do not know. These *Oceanodroma* are evidently referable to three different species, viz.:

1. *O. macrodactyla* W. E. Bryant from Guadalupe Island, a large species with partly white upper tail coverts.

2. *O. melania* (Bp.) from San Benito Island, a large species of nearly uniform blackish coloration, but with whitish brown apical borders to the upper wing coverts. Erroneously labelled as *O. socorroensis*.

3. A species evidently closely allied to *O. monorhis* (Swinh.), but erroneously labelled as "*O. melania*." From the latter the two specimens from San Benito Island now in my collection differ in their much smaller size and in having the front, face and chin suffused with grayish slate color instead of being uniform with the rest of the body, also in the back being paler and of a more brownish black. Otherwise the coloration in these two birds is the same as in *O. melania*.

They are labelled as follows: *Oceanodroma melania*, San Benito Island, Lower California, July 14, 1897. "♂" and "♀" respectively, numbers 1770, 1763.

When at London last year by the kindness of the Hon. Walter Rothschild, I had an opportunity to compare my small so-called *O. melania* with specimens of *O. monorhis* Swinh. from Amuria in the Tring Museum. I found them to be closely allied to *O. monorhis*, but differing from it in their much shorter wings, also somewhat shorter tail and bill, and in their darker and less grayish upper parts.

O. monorhis had not yet been mentioned as occurring in San Benito Island, or elsewhere on the coast of California. The new form as described above is evidently the American representative of that Asiatic species.

While participating in the fourth International Congress at London, I had also the good fortune to show my birds to Mr. Frank M. Chapman of New York. Mr. Chapman having satisfied himself of the distinctness of this new form, I have great pleasure in naming it after a distinguished authority on North American birds.



THE STATUS OF THE "SAN FRANCISCO TITMOUSE."

BY JOSEPH GRINNELL.

TOWARDS the latter part of 1903 a new form of the Plain Titmouse was described¹ under the name *Bæolophus inornatus restrictus*. It was based upon specimens from a suspiciously restricted locality, namely the vicinity of Oakland, California. Without any apparent hitch the A. O. U. Committee on Nomenclature published² its acceptance of the alleged subspecies just nine months

¹ Proc. Biol. Soc. Wash., XVI, Sept. 30, 1903, p. 109.

² Auk, XXI, July, 1904, p. 418.